

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-00-

ed
0188

Public reporting burden for this collection of information is estimated to average 1 hour per record, including gathering and maintaining the data needed, and completing and reviewing the collection of information, including suggestions for reducing the burden, to Washington Headquarters Service, Paperwork Project, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

listing data sources,
other aspect of this
parts, 1215 Jefferson
DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED Final - 15 November 1996 - 14 February 2000	
4. TITLE AND SUBTITLE Reactions of Laser-Ablated Metal Atoms				5. FUNDING NUMBERS F49620-97-1-0010	
6. AUTHOR(S) Dr. Lester Andrews Department of Chemistry					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Virginia Charlottesville, VA 22904-4319				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL 801 North Randolph Street, Room 732 Arlington, VA 22203-1977				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Laser-ablated boron atoms react with methyl fluoride in an argon stream to form two major products, CH ₂ BF and CHBF. In similar reactions of boron with methyl chloride and methyl bromide, both CH ₂ BX and CHBX are also observed, as well as the primary insertion products, CH ₃ BCl and CH ₃ BBr, respectively. Infrared spectra of isotopic combinations of B and CH ₃ X and density functional theory frequency calculations provide evidence for the product identification. Both products form via insertion into the C-X bond followed by loss of one or two H atoms for all reactions. Comparisons are made with the products of the reactions of B with C ₂ H ₆ , CH ₃ NH ₂ , and CH ₃ OH, all of which are isoelectronic with CH ₃ F. The calculations not only predict the vibrational frequencies exceptionally well, but also predict that there is no dative bonding in either CH ₂ BX or CHBX, despite the empty p-orbital on B and the filled p-orbitals on the halogen atom.					
14. SUBJECT TERMS Atoms, metal				15. NUMBER OF PAGES 3	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclass		18. SECURITY CLASSIFICATION OF THIS PAGE Unclass		19. SECURITY CLASSIFICATION OF ABSTRACT Unclass	
20. LIMITATION OF ABSTRACT					

20001016 020

DTIC QUALITY INSPECTED 4

Final Technical Report to the AFOSR

Lester Andrews

Chemistry Department

University of Virginia

Charlottesville, VA 22904-4319

email: lsa@virginia.edu

F49620-97-1-0010

Reactions of Laser-Ablated Metal Atoms

September 1, 1998 to February 14, 2000

ACCOMPLISHMENTS

Research under Air Force support in the final year has involved reactions of laser-ablated metal atoms and hydrogen.

Reactions of laser-ablated ruthenium atoms with carbon monoxide and hydrogen in solid argon produce the unsaturated ruthenium carbonyl dihydride $\text{H}_2\text{Ru}(\text{CO})_2$ and the hydrogen complexes $(\text{H}_2)_x\text{RuCO}$ ($x=1,2$). The observed absorption bands of the reaction products are identified by isotopic substitution and reproduced well by density functional theory calculations of vibrational fundamentals. The growth of bands due to $\text{H}_2\text{Ru}(\text{CO})_2$ and $(\text{H}_2)\text{RuCO}$ during annealing in solid argon indicates that ruthenium monocarbonyl is coordinated with H_2 to form the dihydrogen complex $(\text{H}_2)\text{RuCO}$, while $\text{H}_2\text{Ru}(\text{CO})_2$ is formed from the $(\text{H}_2)\text{RuCO}$ complex inserting into (H_2) upon coordinating with another CO.

Infrared spectra of representative LnH_{1-4} lanthanide hydride molecules are observed in solid argon following reactions of the laser-ablated metal atoms with H_2 . The LnH_2 dihydride is

the major product. Spectra of isolated molecular EuH_2 and solid stoichiometric EuH_2 reveal close correspondence lending support to the molecular assignment, and by extrapolation, providing benchmark infrared frequencies for other lanthanide hydride solids based on their respective molecular vibrational modes.

Laser-ablated boron atoms react with methyl fluoride in an argon stream to form two major products, CH_2BF and CHBF . In similar reactions of boron with methyl chloride and methyl bromide, both CH_2BX and CHBX are also observed, as well as the primary insertion products, CH_3BCl and CH_3BBR , respectively. Infrared spectra of isotopic combinations of B and CH_3X and density functional theory frequency calculations provide evidence for the product identification. Both products form via insertion into the C-X bond followed by loss of one or two H atoms for all reactions. Comparisons are made with the products of the reactions of B with C_2H_6 , CH_3NH_2 and CH_3OH , all of which are isoelectronic with CH_3F . The calculations not only predict the vibrational frequencies exceptionally well, but also predict that there is no dative bonding in either CH_2BX or CHBX , despite the empty p-orbital on B and the filled p-orbitals on the halogen atom.

PERSONNEL SUPPORTED

Xuefeng Wang and Stephen Willson

PUBLICATIONS

1. L. Andrews, P. F. Souter, W. D. Bare, and B. Liang, "Reactions of Laser-Ablated Mo and W Atoms with Dinitrogen: Infrared Spectra of Metal Nitrides, Dinitrides, and Complexes in Solid Argon and Nitrogen," *J. Phys. Chem. A* **1999**, *103*, 4649-4658.
2. L. Andrews, G. P. Kushto, M. F. Zhou, S. P. Willson, and P. F. Souter, "Infrared Spectrum of CCH^+ in Solid Argon and Neon," *J. Chem. Phys.* **1999**, *110*, 4457-4466.

3. S. P. Willson and L. Andrews, "Characterization of the Reaction Products of Laser-Ablated Lanthanide Metal Atoms with Molecular Hydrogen. Infrared Spectra of LnH, LnH₂, LnH₃, and LnH₄ Molecules in Solid Argon," *J. Phys. Chem. A* **2000**, *104*, 1640-1647.
4. X. Wang and L. Andrews, "Reactions of Laser-Ablated Ruthenium Atoms with CO and H₂ Mixtures: Infrared Spectra and Density Functional Theory Calculations of H₂Ru(CO)_x (x=2-4) and (H₂)RuCO," *J. Phys. Chem. A* **2000**, in press.
5. D. V. Lanzisera and L. Andrews, "Reactions of Laser-Ablated Boron Atoms with Methyl Halides in Excess Argon. Infrared Spectra and Density Functional Theory Calculations on CH₃BX, CH₂BX and CHBX (X=F, Cl, Br)," *J. Phys. Chem. A* **2000**, in press.

INTERACTIONS

1. Invited Lecture: "Matrix Isolation – Past, Present, Future," Gordon Research Conference, Chemistry and Physics of Matrix Isolated Species, July 1999.
2. Invited Lecture: "Reactions of Laser-Ablated Transition Metal Atoms, Cations and Electrons with Small Molecules," Twelfth International Conference on Fourier Transform Spectroscopy, Waseda University, Tokyo, Japan, August 1999.
3. Visited Air Force Research Laboratories, host Dr. Mario Fajardo, November 19, 1999.